

Hannes Strass

(based on slides by Michael Thielscher)

Faculty of Computer Science, Institute of Artificial Intelligence, Computational Logic Group


Negation: Model Theory

Lecture 9, 12th Dec 2022 // Foundations of Logic Programming, WS 2022/23

Previously ...

- For every normal logic program P , its **completion** $comp(P)$ replaces the logical implications of clauses by equivalences.
- SLDNF resolution w.r.t. P is **sound** for entailment w.r.t. $comp(P)$.
- SLDNF resolution is only **complete** (for entailment w.r.t. $comp(P)$) for certain combinations of classes of programs, queries, and selection rules.
- For a normal program P , its **dependency graph** D_P explicitly shows positive and negative dependencies between predicate symbols.
- A normal program P is **stratified** iff D_P has no cycle with a negative edge.

$P:$

$$p \leftarrow q, \sim r$$
$$q \leftarrow r$$


Completion of $P:$

$$p \leftrightarrow (q \wedge \neg r)$$

$$q \leftrightarrow r$$

$$r \leftrightarrow \text{false}$$

Overview

Extended Consequence Operator

Standard Models

Perfect Models and Local Stratification

Well-Supported Models

Extended Consequence Operator

Extended Consequence Operator

Definition

Let P be a normal program and I be a Herbrand interpretation. Then

$$T_P(I) := \{H \mid H \leftarrow \vec{B} \in \text{ground}(P), I \models \vec{B}\}$$

In case P is a definite program, we know that

- T_P is monotonic,
- T_P is continuous,
- T_P has the least fixpoint $\mathcal{M}(P)$,
- $\mathcal{M}(P) = T_P \uparrow \omega$.

For normal programs, all of these properties are lost.

Extended T_P -Characterization (1)

Lemma 4.3

Let P be a normal program and I be a Herbrand interpretation. Then

$$I \models P \quad \text{iff} \quad T_P(I) \subseteq I$$

Proof.

$$\begin{aligned} & I \models P \\ \text{iff} & \text{ for every } H \leftarrow \vec{B} \in \text{ground}(P) : I \models \vec{B} \text{ implies } I \models H \\ \text{iff} & \text{ for every } H \leftarrow \vec{B} \in \text{ground}(P) : I \models \vec{B} \text{ implies } H \in I \\ \text{iff} & \text{ for every ground atom } H : H \in T_P(I) \text{ implies } H \in I \\ \text{iff} & T_P(I) \subseteq I \end{aligned}$$

□

Extended T_P -Characterization (2)

Definition

Let F and Π be ranked alphabets of function symbols and predicate symbols, respectively, let $= \in \Pi$ be a binary predicate symbol (**equality**), and let I be a Herbrand interpretation for F and Π . Then

$$I_{=} := I \cup \{=(t,t) \mid t \in HU_F\}$$

is called a **standardized** Herbrand interpretation for F and $\Pi \cup \{=\}$.

Lemma 4.4

Let P be a normal program and I a Herbrand interpretation. Then

$$I_{=} \models \text{comp}(P) \quad \text{iff} \quad T_P(I) = I$$

Extended T_P -Characterization (3)

Proof Idea of Lemma 4.4:

$$I_{=} \models \text{comp}(P)$$

iff (since $I_{=}$ is a model for standard axioms of equality and inequality)

$$\text{for every ground atom } H: I_{=} \models \left(H \leftrightarrow \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B} \right)$$

iff for every ground atom H : $H \in I$ iff $I \models \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B}$

iff for every ground atom H : $H \in I$ iff $I \models \vec{B}$ for some $H \leftarrow \vec{B} \in \text{ground}(P)$

iff for every ground atom H : $H \in I$ iff $H \in T_P(I)$

iff $T_P(I) = I$



Completion may be Inadequate

Consider the following normal logic program P :

$$\begin{aligned}ill &\leftarrow \sim ill, infection \\infection &\leftarrow\end{aligned}$$

Its completion $comp(P) \supseteq \{ill \leftrightarrow (\neg ill \wedge infection), \quad infection \leftrightarrow true\}$ is unsatisfiable (it has no models).

Hence, $comp(P) \not\models healthy$.

But $I = \{infection, ill\}$ is the only Herbrand model of P :

$$\{ill \leftarrow (\neg ill \wedge infection), infection\} \equiv \{ill \vee \neg \neg ill \vee \neg infection, infection\}$$

Hence, $P \not\models healthy$.

Non-Intended Minimal Herbrand Models

$$P_1: \quad p \leftarrow \sim q$$

P_1 has three Herbrand models:

$$M_1 = \{p\}, M_2 = \{q\}, \text{ and } M_3 = \{p, q\}.$$

P_1 has no least, but two minimal Herbrand models: M_1 and M_2

However: M_1 , and not M_2 , is the “intended” model of P_1 .

Supported Herbrand Interpretations

Definition

A Herbrand interpretation I of P is **supported**

$:\Leftrightarrow$ for every $H \in I$ there exists some $H \leftarrow \vec{B} \in \text{ground}(P)$ such that $I \models \vec{B}$.

If additionally $I \models P$, we say that I is a **supported model** of P .

(Intuitively: \vec{B} is an explanation for H .)

Example

- M_1 is a supported model of P_1 . (Literal $\sim q$ is a support for p .)
- M_2 is no supported model of P_1 . (Atom $q \in M_2$ has no support.)
- Note (cf. Lemma 4.3) that $T_{P_1}(M_2) = \emptyset \subsetneq M_2$, but in contrast $T_{P_1}(M_1) = M_1$.
- The definite (therefore normal) program $\{p \leftarrow q, \quad q \leftarrow p\}$ has two supported models: \emptyset and $\{p, q\}$. In the second supported model, p is an explanation for q and vice versa. Thus “support” can be cyclic.

Extended T_P -Characterization (4)

Lemma 6.2

Let P be a normal program and I be a Herbrand interpretation. Then

I is a supported model of P iff $T_P(I) = I$

Proof Idea.

- $I \models P$ and I supported
- iff for every $(H \leftarrow \vec{B}) \in \text{ground}(P)$: $I \models \vec{B}$ implies $I \models H$
and for every $H \in I$: $I \models \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B}$
- iff for every ground atom H : $I \models \left(H \leftarrow \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B} \right)$
and $I \models \left(H \rightarrow \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B} \right)$
- iff for every ground atom H : $I \models \left(H \leftrightarrow \bigvee_{(H \leftarrow \vec{B}) \in \text{ground}(P)} \vec{B} \right)$
- iff $I \models$ model for $\text{comp}(P)$
- iff (Lemma 4.4) $T_P(I) = I$

□

Standard Models

Non-Intended Supported Models

$$P_2: \quad \begin{array}{l} p \leftarrow \sim q \\ q \leftarrow q \end{array}$$

P_2 has three Herbrand models:

$$M_1 = \{p\}, M_2 = \{q\}, \text{ and } M_3 = \{p, q\}$$

P_2 has two supported Herbrand models:

$$M_1 \text{ and } M_2$$

However: M_1 , and not M_2 , is the “intended” model of P_2 .

M_1 will be called the standard model of P_2 (cf. slide 19).

Stratifications

Definition

Let P be a normal program with dependency graph D_P .

- A predicate symbol p is **defined** in P
: $\iff P$ contains a clause $p(t_1, \dots, t_n) \leftarrow \vec{B}$.
- $P_1 \cup \dots \cup P_n = P$ is a **stratification** of P
: \iff
 1. $P_i \neq \emptyset$ for every $i \in [1, n]$
 2. $P_i \cap P_j = \emptyset$ for every $i, j \in [1, n]$ with $i \neq j$
 3. for every p defined in P_i and edge $q \xrightarrow{+} p$ in D_P : q is not defined in $\bigcup_{j=i+1}^n P_j$
 4. for every p defined in P_i and edge $q \xrightarrow{-} p$ in D_P : q is not defined in $\bigcup_{j=i}^n P_j$

Lemma 6.5

A normal program P is stratified iff there exists a stratification of P .

Note: A stratified program may have different stratifications.

Example (1)

The normal logic program P is the following:

```
zero(0) ←  
positive(x) ← num(x), ~zero(x)  
num(0) ←  
num(s(x)) ← num(x)
```

$P_1 \cup P_2 \cup P_3$ is a stratification of P , where

$$P_1 = \{num(0) \leftarrow, num(s(x)) \leftarrow num(x)\}$$

$$P_2 = \{zero(0) \leftarrow\}$$

$$P_3 = \{positive(x) \leftarrow num(x), \sim zero(x)\}$$

Example (1)

The normal logic program P is the following:


```
zero(0) ←  
positive(x) ← num(x), ~zero(x)  
num(0) ←  
num(s(x)) ← num(x)
```

$P_1 \cup P_2 \cup P_3$ is a stratification of P , where

$$P_1 = \{num(0) \leftarrow, num(s(x)) \leftarrow num(x)\}$$

$$P_2 = \{zero(0) \leftarrow\}$$

$$P_3 = \{positive(x) \leftarrow num(x), \sim zero(x)\}$$



Example (2)

$num(0) \leftarrow$
 $num(s(x)) \leftarrow num(x)$
 $even(0) \leftarrow$
 $even(x) \leftarrow \sim odd(x), num(x)$
 $odd(s(x)) \leftarrow even(x)$

P admits no stratification.

Example (2)

$num(0) \leftarrow$
 $num(s(x)) \leftarrow num(x)$
 $even(0) \leftarrow$
 $even(x) \leftarrow \sim odd(x), num(x)$
 $odd(s(x)) \leftarrow even(x)$

P admits no stratification.



Quiz: Stratifications

Quiz

Consider the normal logic program P where x is the only variable: ...

Standard Models (Stratified Programs)

Definition

Let I be an Herbrand interpretation, Π be a set of predicate symbols.

$$I \upharpoonright \Pi := I \cap \{p(t_1, \dots, t_n) \mid p \in \Pi, t_1, \dots, t_n \text{ ground terms}\}$$

Let $P_1 \cup \dots \cup P_n$ be a stratification of the normal program P .

M_1 := least Herbrand model of P_1 such that

$$M_1 \upharpoonright \{p \mid p \text{ not defined in } P\} = \emptyset$$

M_2 := least Herbrand model of P_2 such that

$$M_2 \upharpoonright \{p \mid p \text{ defined nowhere or in } P_1\} = M_1$$

\vdots

M_n := least Herbrand model of P_n such that

$$M_n \upharpoonright \{p \mid p \text{ defined nowhere or in } P_1 \cup \dots \cup P_{n-1}\} = M_{n-1}$$

We call $M_P = M_n$ the **standard model** of P .

Standard Models (Stratified Programs)

Definition

Let I be an Herbrand interpretation, Π be a set of predicate symbols.

$$I \upharpoonright \Pi := I \cap \{p(t_1, \dots, t_n) \mid p \in \Pi, t_1, \dots, t_n \text{ ground terms}\}$$

Let $P_1 \cup \dots \cup P_n$ be a stratification of the normal program P .

M_1 := least Herbrand model of P_1 such that

$$M_1 \upharpoonright \{p \mid p \text{ not defined in } P_1 \cup \dots \cup P_n\} = \emptyset$$

M_2 := least Herbrand model of P_2 such that

$$M_2 \upharpoonright \{p \mid p \text{ not defined in } P_2 \cup \dots \cup P_n\} = M_1$$

\vdots

M_n := least Herbrand model of P_n such that

$$M_n \upharpoonright \{p \mid p \text{ not defined in } P_n\} = M_{n-1}$$

We call $M_P = M_n$ the **standard model** of P .

Example (1)

Let $P_1 \cup P_2 \cup P_3$ with

$$P_1 = \{num(0) \leftarrow, \quad num(s(x)) \leftarrow num(x)\}$$

$$P_2 = \{zero(0) \leftarrow\}$$

$$P_3 = \{positive(x) \leftarrow num(x), \sim zero(x)\}$$

be a stratification of P . Then:

$$\begin{aligned} M_1 &= \{num(t) \mid t \in HU_{\{s,0\}}\} &&= \{num(0), num(s(0)), \dots\} \\ M_2 &= \{num(t) \mid t \in HU_{\{s,0\}}\} \cup \{zero(0)\} &&= \{zero(0), num(0), num(s(0)), \dots\} \\ M_3 &= \{num(t) \mid t \in HU_{\{s,0\}}\} \cup \{zero(0)\} &&= \{zero(0), num(0), num(s(0)), \dots\} \\ &\cup \{positive(t) \mid t \in HU_{\{s,0\}} \setminus \{0\}\} &&\cup \{positive(s(0)), positive(s(s(0))), \dots\} \end{aligned}$$

Hence $M_P = M_3$ is the standard model of P .

Properties of Standard Models

Theorem 6.7

Consider a stratified program P . Then:

- M_P does not depend on the chosen stratification of P ,
- M_P is a minimal model of P ,
- M_P is a supported model of P .

Corollary

For a stratified program P , $comp(P)$ admits a Herbrand model.

Perfect Models and Local Stratification

Stratification may be too demanding

Consider the first-order program P_1 over $\Pi_1 = \{even/1\}$ and $F_1 = \{s/1, 0/0\}$:

- P_1 is **not stratified**, since $even/1$ depends negatively on itself.

```
even(0) ←  
even(s(x)) ← ~even(x)
```

Observation

P_1 has a clear intended model: $\{even(0), even(s(s(0))), even(s(s(s(s(0))))), \dots\}$.

Consider, in contrast, the **propositional** program P_0 over $\Pi_0 = HB_{\{even\}, \{s, 0\}} = \{even(0)/0, even(s(0))/0, \dots\}$ and $F_0 = \emptyset$:

- P_0 is **stratified**.
- The standard model of P_0 is the intended model of P_1 .

```
even(0) ←  
even(s(0)) ← ~even(0)  
even(s(s(0))) ← ~even(s(0))  
⋮
```

Perfect Models

Definition

Let P be a normal program over Π and F , and let \prec be a well-founded ordering on $HB_{\Pi,F}$. Further, let M and N be Herbrand interpretations of P .

- N is **preferable** to M (written $N \triangleleft M$)
: \iff for every $B \in N \setminus M$ there exists an $A \in M \setminus N$ such that $A \prec B$.
- A Herbrand model M of P is **perfect** (w.r.t. \prec)
: \iff there is no Herbrand model of P that is preferable to M .

Well-founded orderings admit no infinite descending chains $\dots \prec c_2 \prec c_1 \prec c_0$.

Example

$$\begin{array}{l} p \leftarrow \sim q \\ q \leftarrow q \end{array}$$

For the well-founded ordering $q \prec p$, we obtain $\{p\} \triangleleft \{q\}$.

The Standard Model is Perfect

Lemma 6.9

Let P be a normal program and $<$ be a well-founded ordering on $HB_{\Pi,F}$.

- If $N \subsetneq M$ then $N < M$.
- Every perfect model of P is minimal.
- The relation $<$ is a partial order on Herbrand interpretations.

Theorem 6.10

Let P be a stratified normal program over Π and F and for $A, B \in HB_{\Pi,F}$ define $A < B : \iff$ the predicate symbol of B depends negatively on the predicate symbol of A .

Then M_P is a unique perfect model of P .

The standard model M_P is thus the $<$ -least Herbrand model of P .

The Standard Model is Perfect

Lemma 6.9

Let P be a normal program and $<$ be a well-founded ordering on $HB_{\Pi,F}$.

- If $N \subsetneq M$ then $N < M$.
- Every perfect model of P is minimal.
- The relation $<$ is a partial order on Herbrand interpretations.

Theorem 6.10

Let P be a stratified normal program over Π and F and for $A, B \in HB_{\Pi,F}$ define $A < B : \iff$ the predicate symbol of B depends negatively on the predicate symbol of A .

Then M_P is a unique perfect model of P .

The standard model M_P is thus the $<$ -least Herbrand model of P .

But how to come up with an ordering $<$ for non-stratified programs?

Local Stratification

Definition

Let P be a normal program over Π and F .

- A **local stratification** for P is a function $strat$ from $HB_{\Pi,F}$ to the countable ordinals.
- For a given local stratification $strat$ and $A \in HB_{\Pi,F}$, we define $strat(\sim A) = strat(A) + 1$.
- A clause $c \in P$ is **locally stratified w.r.t. $strat$**
: \iff for every $A \leftarrow \vec{K}, L, \vec{M} \in ground(c)$, we have $strat(A) \geq strat(L)$.
- P is **locally stratified w.r.t. $strat$**
: \iff all $c \in P$ are locally stratified w.r.t. $strat$.
- P is **locally stratified**
: \iff it is locally stratified w.r.t. to some local stratification.

\rightsquigarrow A first-order program is locally stratified if its ground version is stratified.

Locally Stratified Programs & Perfect Models

Lemma 6.12

Every stratified program is locally stratified.

Example

The program

$$\begin{aligned} \text{even}(0) &\leftarrow \\ \text{even}(s(x)) &\leftarrow \sim\text{even}(x) \end{aligned}$$

is locally stratified, but not stratified.

Theorem 6.13

Let P be a normal logic program (over Π and F) that is locally stratified (w.r.t. strat), and for $A, B \in HB_{\Pi, F}$ define $A < B :\iff \text{strat}(A) < \text{strat}(B)$.

Then P has a unique perfect model.

Well-Supported Models

From Supported to Well-Supported Models

$$p \leftarrow q$$
$$q \leftarrow p$$

has two supported models, \emptyset and $\{p, q\}$.

Only the minimal supported model is intended.

From Supported to Well-Supported Models

$$p \leftarrow q$$
$$q \leftarrow p$$

has two supported models, \emptyset and $\{p, q\}$.

Only the minimal supported model is intended.

$$p \leftarrow \sim q$$
$$q \leftarrow q$$

has two minimal supported models, $\{p\}$ and $\{q\}$.

Only $\{p\}$ is intended: the support of q
("q because q") is unfounded.

From Supported to Well-Supported Models

$$\begin{aligned} p &\leftarrow q \\ q &\leftarrow p \end{aligned}$$

has two supported models, \emptyset and $\{p, q\}$.

Only the minimal supported model is intended.

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow q \end{aligned}$$

has two minimal supported models, $\{p\}$ and $\{q\}$.

Only $\{p\}$ is intended: the support of q (" q because q ") is unfounded.

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow \sim p \end{aligned}$$

has two minimal supported models, $\{p\}$ and $\{q\}$.

- The program is not (locally) stratified.
- The situation is symmetric, so why should we prefer one model over the other?

Well-Supported Models

Definition

Let P be a normal logic program over vocabulary Π, F .

A Herbrand interpretation $I \subseteq HB_{\Pi, F}$ is **well-supported**

$:\Leftrightarrow$

there is a well-founded ordering $<$ on $HB_{\Pi, F}$ such that:

for each $A \in I$ there is a clause $A \leftarrow \vec{B} \in \text{ground}(P)$ with:

- $I \models \vec{B}$
- for every positive atom $C \in \vec{B}$, we have $C < A$.

If additionally $I \models P$, then I is a **well-supported model** of P .

Well-supported models disallow circular justifications.

Theorem 6.20

Any locally stratified normal logic program P has a unique well-supported model that coincides with its perfect model.

Well-Supported Models: Examples

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow q \end{aligned}$$

has $\{p\}$ as only well-supported model.

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow \sim p \end{aligned}$$

has two well-supported models, $\{p\}$ and $\{q\}$.

$$\begin{aligned} p &\leftarrow q \\ p &\leftarrow \sim q \\ q &\leftarrow p \\ q &\leftarrow \sim p \end{aligned}$$

has no well-supported model.

Well-Supported Models: Examples

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow q \end{aligned}$$

has $\{p\}$ as only well-supported model.

$$\begin{aligned} p &\leftarrow \sim q \\ q &\leftarrow \sim p \end{aligned}$$

has two well-supported models, $\{p\}$ and $\{q\}$.

$$\begin{aligned} p &\leftarrow q \\ p &\leftarrow \sim q \\ q &\leftarrow p \\ q &\leftarrow \sim p \end{aligned}$$

has no well-supported model.

Preview: Well-supported models are also known as **stable models**.

Conclusion

Summary

- The immediate consequence operator T_P for a normal logic program P characterizes the **supported models** of P (= the models of $comp(P)$).
- The **stratification** of a program P partitions the program in layers (strata) such that predicates in one layer only **negatively/positively** depend on predicates in **strictly lower/lower or equal** layers.
- Programs P that are **stratified** have an intended **standard model** M_P .
- A program is **locally stratified** iff its “propositional version” is stratified.
- Locally stratified programs allow for a unique **perfect model**.
- A normal program P may have zero or more **well-supported models**.

Suggested action points:

- Prove Lemma 6.5; show that every well-supported model is supported.